



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

Widespread anthelmintic resistance in European farmed ruminants: a systematic review

Rose, H ; Rinaldi, L ; Bosco, A ; Mavrot, F ; de Waal, T ; Skuce, P ; Charlier, J ; Torgerson, P R ;
Hertzberg, H ; Hendrickx, G ; Vercruysse, J ; Morgan, E R

DOI: <https://doi.org/10.1136/vr.102982>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-110037>

Journal Article

Accepted Version

Originally published at:

Rose, H; Rinaldi, L; Bosco, A; Mavrot, F; de Waal, T; Skuce, P; Charlier, J; Torgerson, P R; Hertzberg, H; Hendrickx, G; Vercruysse, J; Morgan, E R (2015). Widespread anthelmintic resistance in European farmed ruminants: a systematic review. *Veterinary Record*, 176(21):546.

DOI: <https://doi.org/10.1136/vr.102982>

Widespread anthelmintic resistance in European farmed ruminants: a systematic review

Rose, H.^{a,b} (BSc, PhD), Rinaldi, L.^c (BSc, PhD, AssEVPC), Bosco, A.^c (DVM, PhD), Mavrot, F.^d (DVM), de Waal, T.^e (BVSc, PhD, DipDatMet, HDipUTL, DipEVPC), Skuce, P.^f (BSc, PhD), Charlier, J.^g (DVM, PhD, DipEVPC), Torgerson, P. R.^d (BA, VetMB, PhD, DEVPH, MRCVS), Hertzberg, H.^h (DVM, DipEVPC), Hendrickx, G.ⁱ (DVM, PhD), Vercruysse, J.^g (DVM, DipEVPC), Morgan, E. R.^{b,j} (MA, VetMB, PhD, DipEVPC, MRCVS)

^aSchool of Biological Sciences, University of Bristol, Tyndall Avenue, Bristol BS8 1TQ, UK

^bCabot Institute, University of Bristol, Cantocks Close, Bristol BS8 1TS, UK

^cDepartment of Veterinary Medicine and Animal Productions, University of Naples Federico II, CREMOPAR, Regione Campania, Italy

^dSection of Veterinary Epidemiology, University of Zurich, Winterthurerstrasse 270, CH-8057 Zurich, Switzerland

^eSchool of Veterinary Medicine, University College Dublin, Belfield, Dublin, Ireland

^fMoredun Research Institute, Pentlands Science Park, Midlothian EH26 0PZ, UK

^gDepartment of Virology, Parasitology and Immunology, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, 9820 Merelbeke, Belgium

^hInstitute of Parasitology, University of Zurich, Winterthurerstrasse 266a, CH-8057 Zurich, Switzerland

ⁱAvia-GIS BVBA, Risschotlei 33, 2980 Zoersel, Belgium

^jSchool of Veterinary Sciences, University of Bristol, Langford House, Bristol BS40 5DU, UK

E-mail for correspondence: hannah.rose@bristol.ac.uk

Anthelmintic resistance (AR) in gastrointestinal nematodes (GINs) has been reported worldwide in multiple nematode and livestock species (Kaplan and Vidyashankar 2012) and is a major constraint on production on affected farms (Sutherland and others 2010, Miller and others 2012). In the UK and Ireland for example, AR in GINs and anthelmintic treatment failure is widespread in sheep (e.g. Bartley and others 2003, Keane and others 2014) and increasingly reported in cattle (e.g. O'Shaughnessy and others 2014). There is, therefore, a need to develop and adopt GIN control strategies that maintain the efficacy of anthelmintics and to identify risk factors for the development of AR.

Environmental constraints on farm management and the survival of nematodes *in refugia* appear to play an important role in the development of AR. In a random survey of sheep farms in Norway, AR was found only in coastal regions (Domke and others 2012a). Papadopoulos and others (2001) observed a higher incidence of AR on isolated Greek islands, suggesting that drought hastens the development of AR. In contrast, Rinaldi and others (2014) observed high anthelmintic efficacy in sheep in southern Italy, despite the Mediterranean climate. This was attributed to the low number of anthelmintic treatments (usually 2 per year) and absence of anthelmintic treatments during periods of drought, when environmental constraints on the free-living stages are highest. Calvete and others (2012) identified an association between AR, distance between farms with AR, management and bioclimatic variables on sheep farms in Aragon, Spain. In particular, the association between AR and climatic conditions was attributed to the application of anthelmintic treatments during the winter months, which increases the selection pressure on the already depleted population of nematodes *in refugia*. Such spatial analyses provide useful insights into risk factors for AR but their application is likely to be limited outside of the region studied. Pan-European spatial analysis and modelling of the distribution of AR may enable the elucidation of common risk factors for the development of AR in European livestock.

A systematic review of peer-reviewed literature was undertaken to record the current distribution of AR in the major GINs (*Teladorsagia* spp., *Trichostrongylus* spp., *Haemonchus contortus*, *Ostertagia ostertagi* and *Cooperia oncophora*) infecting goats, sheep and cattle in Europe (defined as the

European Union (EU), European Economic Area (EEA) and Switzerland). The ISI Web of Science database was explored using the keywords “anthelmintic resistance” (last searched 02/10/14). No restrictions were placed on publication dates. The search yielded 1,852 publications, of which 120 publications were selected based on title and abstract, excluding studies on non-ovine, -bovine or -caprine hosts and nematodes, non-European studies and studies where AR arose through artificial selection. A further nine reports of AR were identified from citations, MSc/PhD theses and authors’ unpublished data. Of these publications, 73 provided reports of AR in cattle, sheep or goats assessed in accordance with World Association for the Advancement of Veterinary Parasitology guidelines (Coles and others 1992) and stated the country or region where the farms were located.

AR in GINs, assessed primarily using faecal egg count reduction tests (FECRT), is widespread in Europe (supplementary figure). Overall, AR was reported in all five GIN genera and in 16 countries throughout Europe (supplementary figure and table). Multiple drug resistance (MDR) in the three main GIN genera infecting sheep and goats was reported in 10 countries (supplementary table). Not all studies tested multiple anthelmintics and, therefore, MDR is likely to be more widespread. Monepantel resistance was reported on sheep farms in the Netherlands in November 2014 (Anon 2014) but was not included in the systematic review as details regarding the methods used to assess resistance were not available at the time of writing. AR against derquantel had not been reported in Europe at the time of writing. However, due to publication and sample selection bias, the absence of reports of AR in some regions may simply be due to a lack of monitoring and AR cannot be considered absent elsewhere. Heterogeneity in the distribution of AR in Europe might also depend on the lack of standardized procedures for surveys and detection of AR on farms and in laboratories.

The estimated prevalence of AR varied by region, anthelmintic class and host. Random surveys of sheep farms have detected albendazole resistance on 11% of farms in Norway (n=19; Domke and others 2012a), ivermectin, benzimidazole and levamisole resistance on 23%, 3.7% and 7.4% of farms, respectively, in Slovakia (n=27; Čerňanská and others 2006), and benzimidazole and levamisole resistance on 83% and 50%, respectively, of farms in Western France (n=23; Chartier and others 1998). In the latter study, benzimidazole resistance was also detected on 93% of goat farms (n=15). A further random survey of dairy goat farms in Southwestern France detected benzimidazole resistance on 83% of farms, and multiple resistance to benzimidazole and levamisole on 11% of farms (n=18; Chartier and others 2001). A sample size-weighted mean prevalence of benzimidazole resistance in GINs in sheep and goats of 50.1% was estimated from the above four studies. Excluding goats, the sample size-weighted mean prevalence of benzimidazole resistance in sheep GINs was 32.1%. Insufficient data were available to estimate mean prevalence for other anthelmintic classes and cattle. The prevalence of AR has also been estimated elsewhere e.g. treatment failure has been identified on 51% of Irish sheep farms surveyed (Keane and others 2014) and 64% of Scottish sheep farms surveyed (Bartley and others 2003). These studies provide valuable prevalence estimates but differences in sample (farm) selection methods introduce potential sample selection bias and may affect estimates and comparability between regions. For example, Domke and others (2012a) observed AR on 33% of randomly selected sheep flocks and 80% of non-randomly selected sheep flocks in the Rogaland region of Norway. Therefore, it is recommended that future prevalence surveys follow a random or stratified sampling approach where possible to reduce sample selection bias.

The biases described above currently prevent robust spatial meta-analysis of AR in Europe and restrict the spatial analysis that can be undertaken. In addition, since spatial analysis is rarely the purpose of a study into AR and due to data protection responsibilities, cases are usually reported at a country or regional level. Due to the significant within-region heterogeneity in the distribution of AR (e.g. Calvete and others 2012), data with a higher spatial resolution are required.

Taken together, the peer-reviewed literature paints a picture of widespread AR in Europe with the potential for high regional prevalence.. Veterinarians should continue to promote sustainable anthelmintic use (e.g. Abbott and others 2012, Charlier and others 2014), even on farms where AR is not suspected. Continued surveillance of AR in Europe, reporting the absence of resistance (Paraud and others 2010, Rinaldi and others 2014) and reporting cases in a way that enables spatial meta-analysis will aid in the future identification of risk factors and evaluation of sustainable nematode control practices.

Acknowledgements

The work was supported by funding from the FP7 GLOWORM project – Grant agreement N° 288975CP-TP-KBBE.2011.1.3-04

References

- ANON (2014) Monepantel resistance reported on Dutch sheep farms. *Veterinary Record* 175, 418
- ABBOTT, K. A., TAYLOR, M. A. & STUBBINGS, L. A. (2012) Sustainable worm control strategies for sheep: A technical manual for veterinary surgeons and advisers, 4th Edition.
<http://www.scops.org.uk/content/SCOPS-Technical-manual-4th-Edition-June-2012.pdf>. Accessed July 17, 2014
- ALVAREZ-SÁNCHEZ, M. A., PÉREZ-GARCÍA, J., CRUZ-ROJO, M. A. & ROJO-VÁZQUEZ, F. A. (2006) Anthelmintic resistance in trichostrongylid nematodes of sheep farms in Northwest Spain. *Parasitology Research* 99, 78–83
- ARESKOG, M., LJUNGSTRÖM, B. & HÖGLUND, J. (2013) Limited efficacy of pour-on anthelmintic treatment of cattle under Swedish field conditions. *International Journal for Parasitology: Drugs and Drug Resistance* 3, 129–34
- ARTHO, R., SCHNYDER, M., KOHLER, L., TORGERSON, P. R. & HERTZBERG, H. (2007) Avermectin-resistance in gastrointestinal nematodes of Boer goats and Dorper sheep in Switzerland. *Veterinary Parasitology* 144, 68–73
- BARTLEY, D. J., DONNAN, A. A., JACKSON, E., SARGISON, N., MITCHELL, G. B. B. & JACKSON, F. (2006) A small scale survey of ivermectin resistance in sheep nematodes using the faecal egg count reduction test on samples collected from Scottish sheep. *Veterinary Parasitology* 137, 112–118
- BARTLEY, D. J., JACKSON, E., JOHNSTON, K., COOP, R. L., MITCHELL, G. B. B., SALES, J. & JACKSON, F. (2003) A survey of anthelmintic resistant nematode parasites in Scottish sheep flocks. *Veterinary Parasitology* 117, 61–71
- BARTLEY, D. J., JACKSON, F., JACKSON, E. & SARGISON, N. (2004) Characterisation of two triple resistant field isolates of *Teladorsagia* from Scottish lowland sheep farms. *Veterinary Parasitology* 123, 189–199
- BARTLEY, D. J., MCARTHUR, C. L., DEVIN, L. M., SUTRA, J. F., MORRISON, A. A., LESPINE, A. & MATTHEWS, J. B. (2012) Characterisation of macrocyclic lactone resistance in two field-derived isolates of *Cooperia oncophora*. *Veterinary Parasitology* 190, 454–460
- BJØRN, H., MONRAD, J. & NANSEN, P. (1991) Anthelmintic resistance in nematode parasites of sheep in Denmark with special emphasis on levamisole resistance in *Ostertagia circumcincta*. *Acta Veterinaria Scandinavica* 32, 145–154
- BOERSEMA, J. H., BORGSTEEDE, F. H., EYSKER, M., HENDRIKX, W. M., JANSEN, J. & SMITH-BUYS, C. M. (1987) Prevalence of benzimidazole resistance of nematodes in sheep in The Netherlands. *Research in Veterinary Science* 43, 18–21
- BORGSTEEDE, F. H. (1986) Resistance of *Cooperia curticei* against fenbendazole. *Research in Veterinary Science* 41, 423–424

141 BORGSTEEDE, F. H. & DUYN, S. P. (1989) Lack of reversion of a benzimidazole resistant strain of
 142 *Haemonchus contortus* after six years of levamisole usage. *Research in Veterinary Science* 47, 270–
 143 272

144 BORGSTEEDE, F. H. M., DERCKSEN, D. D. & HUIJBERS, R. (2007) Doramectin and albendazole
 145 resistance in sheep in The Netherlands. *Veterinary Parasitology* 144, 180–183

146 BORGSTEEDE, F. H. M., GEERTS, S., DE DEKEN, R., KUMAR, V. & BRANDT, J. (1992) Studies on an
 147 *Ostertagia ostertagi* strain suspected to be resistant to benzimidazoles. *Veterinary Parasitology* 41,
 148 85–92

149 BORGSTEEDE, F. H. M., PEKELDER, J. J. & DERCKSEN, D. P. (1996) Anthelmintic resistant nematodes
 150 in goats in The Netherlands. *Veterinary Parasitology* 65, 83–87

151 BORGSTEEDE, F. H. M., PEKELDER, J. J., DERCKSEN, D. P., SOL, J., VELLEMA, P., GAASENBEEK, C. P. &
 152 VAN DER LINDEN, J. N. (1997) A survey of anthelmintic resistance in nematodes of sheep in The
 153 Netherlands. *The Veterinary Quarterly* 19, 167–172

154 BRITT, D. P. & OAKLEY, G. A. (1986) Anthelmintic evaluation of a thiabendazole-resistant strain of
 155 *Ostertagia circumcincta* recovered from sheep in England. *Veterinary Parasitology*, 19, 95–101

156 BURGESS, C. G. S., BARTLEY, Y., REDMAN, E., SKUCE, P. J., NATH, M., WHITELAW, F., TATE, A.,
 157 GILLEARD, J. S. & JACKSON, F. (2012) A survey of the trichostrongylid nematode species present on
 158 UK sheep farms and associated anthelmintic control practices. *Veterinary Parasitology* 189, 299–307

159 CABARET, J., BAUDET, H. M., DEVOS, J., HUBERT, J., CORTET, J. & SAUVÉ, C. (1995) Studies on
 160 mutlispecific resistance of gastrointestinal nematodes to benzimidazoles on dairy-goat farms.
 161 *Veterinary Parasitology* 60, 331–337

162 CALVETE, C., CALAVIA, R., FERRER, L. M., RAMOS, J. J., LACASTA, D. & URIARTE, J. (2012)
 163 Management and environmental factors related to benzimidazole resistance in sheep nematodes in
 164 Northeast Spain. *Veterinary Parasitology* 184, 193–203

165 CAWTHORNE, R. J. & CHEONG, F. H. (1984) Prevalence of anthelmintic resistant nematodes in sheep
 166 in south-east England. *Veterinary Record* 114, 562–564

167 CAWTHORNE, R. J. & WHITEHEAD, J. D. (1983) Isolation of benzimidazole resistant strains of
 168 *Ostertagia circumcincta* from British sheep. *Veterinary Record* 112, 274–277

169 ČERNÁNSKÁ, D., VÁRADY, M. & ČORBA, J. (2006) A survey on anthelmintic resistance in nematode
 170 parasites of sheep in the Slovak Republic. *Veterinary Parasitology* 135, 39–45

171 CHARLIER, J., MORGAN, E. R., RINALDI, L., VAN DIJK, J., DEMELER, J., HÖGLUND, J., HERTZBERG, H.,
 172 VAN RANST, B., HENDRICKX, G., VERCRUYSE, J. & KENYON, F. (2014) Practices to optimise
 173 gastrointestinal nematode control on sheep, goat and cattle farms in Europe using targeted
 174 (selective) treatments. *Veterinary Record* 175, 250–255

175 CHARTIER, C., PORS, I., HUBERT, J., ROCHETEAU, D., BENOIT, C. & BERNARD, N. (1998) Prevalence of
176 anthelmintic resistant nematodes in sheep and goats in Western France. *Small Ruminant Research*
177 29, 33–41

178 CHARTIER, C., SOUBIRAC, F., PORS, I., SILVESTRE, A., HUBERT, J., COUQUET, C. & CABARET, J. (2001)
179 Prevalence of anthelmintic resistance in gastrointestinal nematodes of dairy goats under extensive
180 management conditions in southwestern France. *Journal of Helminthology* 75, 325–330

181 COLES, G. C., BAUER, C., BORGSTEEDE, F. H. M., GEERTS, S., KLEI, T. R., TAYLOR, M. A. & WALLER, P. J.
182 (1992) World Association for the Advancement of Veterinary Parasitology (W.A.A.V.P.) methods for
183 the detection of anthelmintic resistance in nematodes of veterinary importance. *Veterinary*
184 *Parasitology* 44, 35–44

185 COLES, G. C., STAFFORD, K. A. & MACKAY, P. H. (1998) Ivermectin-resistant *Cooperia* species from
186 calves on a farm in Somerset. *Veterinary Record* 142, 255–256

187 CRINGOLI, G., VENEZIANO, V., RINALDI, L., SAUVÉ, C., RUBINO, R., FEDELE, V. & CABARET, J. (2007)
188 Resistance of trichostrongyles to benzimidazoles in Italy: a first report in a goat farm with multiple
189 and repeated introductions. *Parasitology Research* 101, 577–581

190 DEMELER, J., VAN ZEVEREN, A. M. J., KLEINSCHMIDT, N., VERCruysse, J., HÖGLUND, J., KOOPMANN,
191 R., CABARET, J., CLAEREBOU, E., ARESKOG, M. & VON SAMSON-HIMMELSTJERNA, G. (2009)
192 Monitoring the efficacy of ivermectin and albendazole against gastro intestinal nematodes of cattle
193 in Northern Europe. *Veterinary Parasitology* 160, 109–115

194 DÍEZ-BAÑOS, P., PEDREIRA, J., SÁNCHEZ-ANDRADE, R., FRANCISCO, I., SUÁREZ, J. L., DÍAZ, P.,
195 PANADERO, R., ARIAS, M., PAINCEIRA, A., PAZ-SILVA, A. & MORRONGO, P. (2008) Field evaluation for
196 anthelmintic-resistant ovine gastrointestinal nematodes by in vitro and in vivo assays. *The Journal of*
197 *Parasitology* 94, 925–928

198 DOMKE, A. V. M., CHARTIER, C., GJERDE, B., HÖGLUND, J., LEINE, N., VATN, S. & STUEN, S. (2012a)
199 Prevalence of anthelmintic resistance in gastrointestinal nematodes of sheep and goats in Norway.
200 *Parasitology Research* 111, 185–193

201 DOMKE, A. V. M., CHARTIER, C., GJERDE, B. & STUEN, S. (2012b) Benzimidazole resistance of sheep
202 nematodes in Norway confirmed through controlled efficacy test. *Acta Veterinaria Scandinavica* 54,
203 48

204 EL-ABDELLATI, A., CHARLIER, J., GELDHOF, P., LEVECKE, B., DEMELER, J., VON SAMSON-
205 HIMMELSTJERNA, G., CLAEREBOU, E. & VERCruysse, J. (2010a) The use of a simplified faecal egg
206 count reduction test for assessing anthelmintic efficacy on Belgian and German cattle farms.
207 *Veterinary Parasitology* 169, 352–357

208 EL-ABDELLATI, A., GELDHOF, P., CLAEREBOU, E., VERCruysse, J. & CHARLIER, J. (2010b) Monitoring
209 macrocyclic lactone resistance in *Cooperia oncophora* on a Belgian cattle farm during four
210 consecutive years. *Veterinary Parasitology* 171, 167–171

211 GALLIDIS, E., ANGELOPOULOU, K. & PAPADOPOULOS, E. (2012) First identification of benzimidazole
212 resistant *Haemonchus contortus* in sheep in Greece. *Small Ruminant Research* 106, 27–29

213 GEURDEN, T., HOSTE, H., JACQUIET, P., TRAVERSA, D., SOTIRAKI, S., FRANGIPANE DI REGALBONO, A.
 214 and others (2014) Anthelmintic resistance and multidrug resistance in sheep gastro-intestinal
 215 nematodes in France, Greece and Italy. *Veterinary Parasitology* 201, 59–66

216 GOOD, B., HANRAHAN, J. P., DE WAAL, D. T., PATTEN, T., KINSELLA, A. & LYNCH, C. O. (2012)
 217 Anthelmintic-resistant nematodes in Irish commercial sheep flocks-the state of play. *Irish Veterinary*
 218 *Journal* 65, 21

219 GRIMSHAW, W. T., HUNT, K. R., HONG, C. & COLES, G. C. (1994) Detection of anthelmintic resistant
 220 nematodes in sheep in southern England by a faecal egg count reduction test. *Veterinary Record*
 221 135, 372–374

222 HERTZBERG, H., ROSSMANN, J., KOHLER, L. & WILLI, U. (2000) Vorkommen von Benzimidazol-
 223 Resistenzen bei Magen-Darmnematoden des Schafes und der Ziege in der Schweiz. *Wiener*
 224 *Tierärztliche Monatsschrift* 87, 3–9

225 HÖGLUND, J., GUSTAFSSON, K., LJUNGSTRÖM, B.-L., ENGSTRÖM, A., DONNAN, A. & SKUCE, P. (2009)
 226 Anthelmintic resistance in Swedish sheep flocks based on a comparison of the results from the faecal
 227 egg count reduction test and resistant allele frequencies of the beta-tubulin gene. *Veterinary*
 228 *Parasitology* 161, 60–68

229 HONG, C., HUNT, K. R. & COLES, G. C. (1996) Occurrence of anthelmintic resistant nematodes on
 230 sheep farms in England and goat farms in England and Wales. *Veterinary Record* 139, 83–86

231 HONG, C., HUNT, K. R., HARRIS, T. J., COLES, G. C., GRIMSHAW, W. T. & MCMULLIN, P. F. (1992) A
 232 survey of benzimidazole resistant nematodes in sheep in three countries of southern England.
 233 *Veterinary Record* 131, 5–7

234 JACKSON, F., JACKSON, E. & COOP, R. L. (1992) Evidence of multiple anthelmintic resistance in a
 235 strain of *Teladorsagia circumcincta* (*Ostertagia circumcincta*) isolated from goats in Scotland.
 236 *Research in Veterinary Science* 53, 371–374

237 JORDI, R. (1980) Untersuchungen zur Anthelminthika-Resistenz von Trichostrongyliden des Schafes.
 238 *Schweizer Archiv für Tierheilkunde* 122, 679-694

239 KAPLAN, R. M. & VIDYASHANKAR, A. N. (2012) An inconvenient truth: Global worming and
 240 anthelmintic resistance. *Veterinary Parasitology* 186, 70-78

241 KEANE, O. M., KEEGAN, J. D., GOOD, B., DE WAAL, T., FANNING, J., GOTTSTEIN, M., CASEY, M.,
 242 HURLEY, C. & SHEEHAN, M. (2014) High level of treatment failure with commonly used anthelmintics
 243 on Irish sheep farms. *Irish Veterinary Journal* 67, 16

244 MAINGI, N., BJØRN, H., THAMSBORG, S. M., BØGH, H. O. & NANSEN, P. (1996a) A survey of
 245 anthelmintic resistance in nematode parasites of goats in Denmark. *Veterinary Parasitology* 66, 53–
 246 66

247 MAINGI, N., BJØRN, H., THAMSBORG, S. M., BØGH, H. O. & NANSEN, P. (1996b) Anthelmintic
 248 resistance in nematode parasites of sheep in Denmark. *Small Ruminant Research* 23, 171–181

249 MARTÍNEZ-VALLADARES, M., DONNAN, A., GELDHOF, P., JACKSON, F., ROJO-VÁZQUEZ, F.-A. &
 250 SKUCE, P. (2012a) Pyrosequencing analysis of the beta-tubulin gene in Spanish *Teladorsagia*
 251 *circumcincta* field isolates. *Veterinary Parasitology* 184, 371–376

252 MARTÍNEZ-VALLADARES, M., FAMULARO, M. R., FERNÁNDEZ-PATO, N., CORDERO-PÉREZ, C.,
 253 CASTAÑÓN-ORDÓÑEZ, L. & ROJO-VÁZQUEZ, F. A. (2012b) Characterization of a multidrug resistant
 254 *Teladorsagia circumcincta* isolate from Spain. *Parasitology Research* 110, 2083–2087

255 MARTÍNEZ-VALLADARES, M., MARTÍNEZ-PÉREZ, J. M., ROBLES-PÉREZ, D., CORDERO-PÉREZ, C.,
 256 FAMULARO, M. R., FERNÁNDEZ-PATO, N., CASTAÑÓN-ORDÓÑEZ, L. & ROJO-VÁZQUEZ, F. A. (2013)
 257 The present status of anthelmintic resistance in gastrointestinal nematode infections of sheep in the
 258 northwest of Spain by in vivo and in vitro techniques. *Veterinary Parasitology* 191, 177–181

259 MCARTHUR, C. L., BARTLEY, D. J., SHAW, D. J. & MATTHEWS, J. B. (2011) Assessment of ivermectin
 260 efficacy against gastrointestinal nematodes in cattle on four Scottish farms. *Veterinary Record*, 169,
 261 658

262 MCMAHON, C., BARTLEY, D. J., EDGAR, H. W. J., ELLISON, S. E., BARLEY, J. P., MALONE, F. E., HANNA,
 263 RE. E. B., BRENNAN, G. P. & FAIRWEATHER, I. (2013) Anthelmintic resistance in Northern Ireland (I):
 264 prevalence of resistance in ovine gastrointestinal nematodes, as determined through faecal egg
 265 count reduction testing. *Veterinary Parasitology* 195, 122–130

266 MEYER, A. (2001) Verbreitung von benzimidazol-resistenzen bei den trichostrongyliden von schafen
 267 und ziegen in der Schweiz. PhD thesis, University of Zurich

268 MILLER, C., WAGHORN, T., LEATHWICK, D. M., CANDY, P. M., OLIVER, A-M. B. & WATSON, T. G.
 269 (2012) The production cost of anthelmintic resistance in lambs. *Veterinary Parasitology* 186, 376-381

270 MITCHELL, E. S. E., HUNT, K. R., WOOD, R. & MCLEAN, B. (2010) Anthelmintic resistance on sheep
 271 farms in Wales. *Veterinary Record* 166, 650–652

272 MURRI, S., KNUBBEN-SCHWEIZER, G., TORGERSON, P. & HERTZBERG, H. (2014) Frequency of
 273 eprinomectin resistance in gastrointestinal nematodes of goats in canton Berne, Switzerland.
 274 *Veterinary Parasitology* 203, 114–119

275 O'SHAUGHNESSY, J., EARLEY, B., MEE, J. F., DOHERTY, M. L., CROSSON, P., BARRETT, D.,
 276 PRENDIVILLE, R., MACRELLI, M. & DE WAAL, T. (2014) Detection of anthelmintic resistance on two
 277 Irish beef research farms. *Veterinary Record* 175, 120

278 PALCY, C., SILVESTRE, A., SAUVE, C., CORTET, J. & CABARET, J. (2010) Benzimidazole resistance in
 279 *Trichostrongylus axei* in sheep: long-term monitoring of affected sheep and genotypic evaluation of
 280 the parasite. *The Veterinary Journal* 183, 68–74

281 PAPADOPOULOS, E., HIMONAS, C. & COLES, G. C. (2001) Drought and flock isolation may enhance
 282 the development of anthelmintic resistance in nematodes. *Veterinary Parasitology* 97, 253-259

283 PARAUD, C., KULO, A., PORS, I. & CHARTIER, C. (2009) Resistance of goat nematodes to multiple
 284 anthelmintics on a farm in France. *Veterinary Record* 164, 563–564

285 PARAUD, C., PORS, I., REHBY, L. & CHARTIER, C. (2010) Absence of ivermectin resistance in a survey
 286 on dairy goat nematodes in France. *Parasitology Research* 106, 1475-1479

287 PRASLIČKA, J., VÁRADY, M., ČORBA, J. & VESELÝ, L. (1994) A survey of anthelmintic resistance in
 288 Slovakia. *Veterinary Parasitology* 52, 169–171

289 REQUEJO-FERNÁNDEZ, J. A., MARTÍNEZ, A., MEANA, A., ROJO-VÁZQUEZ, F. A., OSORO, K. & ORTEGA-
 290 MORA, L. M. (1997) Anthelmintic resistance in nematode parasites from goats in Spain. *Veterinary*
 291 *Parasitology* 73, 83–88

292 RINALDI, L., MORGAN, E. R., BOSCO, A., COLES, G. C. & CRINGOLI, G. (2014) The maintenance of
 293 anthelmintic efficacy in sheep in a Mediterranean climate. *Veterinary Parasitology* 203, 139-143

294 SARGISON, N. D., JACKSON, F., BARTLEY, D. J. & MOIR, A. C. P. (2005) Failure of moxidectin to control
 295 benzimidazole-, levamisole- and ivermectin-resistant *Teladorsagia circumcincta* in a sheep flock.
 296 *Veterinary Record* 156, 105–109

297 SARGISON, N. D., JACKSON, F., BARTLEY, D. J., WILSON, D. J., STENHOUSE, L. J. & PENNY, C. D. (2007)
 298 Observations on the emergence of multiple anthelmintic resistance in sheep flocks in the south-east
 299 of Scotland. *Veterinary Parasitology* 145, 65–76

300 SARGISON, N. D., SCOTT, P. R., WILSON, D. J., MACRAE, A. I. & PENNY, C. D. (2010) *Teladorsagia*
 301 *circumcincta* resistance to moxidectin and multiple anthelmintic groups in ewes following use of the
 302 persistent drug before lambing. *Veterinary Record* 167, 523–527

303 SCHEUERLE, M. C., MAHLING, M. & PFISTER, K. (2009) Anthelmintic resistance of *Haemonchus*
 304 *contortus* in small ruminants in Switzerland and Southern Germany. *Wiener Klinische Wochenschrift*
 305 121, 46–49

306 SCHNYDER, M., TORGERSON, P. R., SCHÖNMANN, M., KOHLER, L. & HERTZBERG, H. (2005) Multiple
 307 anthelmintic resistance in *Haemonchus contortus* isolated from South African Boer goats in
 308 Switzerland. *Veterinary Parasitology* 128, 285–290

309 STAFFORD, K. & COLES, G. C. (1999) Nematode control practices and anthelmintic resistance in dairy
 310 calves in the south west of England. *Veterinary Record* 144, 659–661

311 SUTHERLAND, A. A., SHAW, J. & SHAW, R. J. (2010) The production costs of anthelmintic resistance
 312 in sheep managed within a monthly preventive drench program. *Veterinary Parasitology* 171, 300-
 313 304

314 TAYLOR, M. A., LEARMOUNT, J., LUNN, E., MORGAN, C. & CRAIG, B. H. (2009) Multiple resistance to
 315 anthelmintics in sheep nematodes and comparison of methods used for their detection. *Small*
 316 *Ruminant Research* 86, 67–70

317 TRAVERSA, D., PAOLETTI, B., OTRANTO, D. & MILLER, J. (2007) First report of multiple drug
 318 resistance in trichostrongyles affecting sheep under field conditions in Italy. *Parasitology Research*
 319 101, 1713–1716

320 VADLEJCH, J., KOPECKÝ, O., KUDRNÁČOVÁ, M., ČADKOVÁ, Z., JANKOVSKÁ, I. & LANGROVÁ, I. (2014)
 321 The effect of risk factors of sheep flock management practices on the development of anthelmintic
 322 resistance in the Czech Republic. *Small Ruminant Research* 117, 183–190

323 VAN DEN BROM, R., MOLL, L., BORGSTEEDE, F. H. M., VAN DOORN, D. C. K., LIEVAART-PETERSON, K.,
 324 DERCKSEN, D. P. & VELLEMA, P. (2013) Multiple anthelmintic resistance of *Haemonchus contortus*,
 325 including a case of moxidectin resistance, in a Dutch sheep flock. *Veterinary Record* 173, 552

326 VÁRADY, M., ČERŇANSKÁ, D. & ČORBA, J. (2006) Use of two in vitro methods for the detection of
 327 anthelmintic resistant nematode parasites on Slovak sheep farms. *Veterinary Parasitology* 135, 325–
 328 331


329 VOIGT, K., SCHEUERLE, M. & HAMEL, D. (2012) Triple anthelmintic resistance in *Trichostrongylus* spp.
 330 in a German sheep flock. *Small Ruminant Research* 106, 30–32

331 WILSON, D. & SARGISON, N. (2007) Anthelmintic resistance in *Teladorsagia circumcincta* in sheep in
 332 the UK. *Veterinary Record* 161, 535–536

333 ZANZANI, S. A., GAZZONIS, A. L., DI CERBO, A., VARADY, M. & MANFREDI, M. T. (2014)
 334 Gastrointestinal nematodes of dairy goats, anthelmintic resistance and practices of parasite control
 335 in Northern Italy. *BMC Veterinary Research* 10, 114

336

Supplementary figure. The distribution of reported cases of anthelmintic resistance (AR) in the European Union, European Economic Area and Switzerland, at national level, based on the systematic review. Shaded countries had at least one reported case of AR. Note that regional distribution within countries is not plotted: see supplementary table for details of drug classes and GIN genera/species, and individual references for specific locations and apparent prevalence. No data were available for Iceland (not shown).



345 **Supplementary table.** Summary of anthelmintic resistance reported in five major gastrointestinal
 346 nematode species/genera infecting goats, sheep and cattle in Europe.

AH class ^a	Host	Region	References
<i>Teladorsagia</i> spp.			
BZ	Sheep	Czech Republic, Denmark, France, Greece, Ireland, Italy, Netherlands, Norway, Slovakia, Spain, UK	Alvarez-Sánchez and others 2006, Bartley and others 2004, 2006, Bjørn and others 1991, Boersema and others 1987, Borgsteede and others 1997, 2007, Britt & Oakley 1986, Cawthorne and Whitehead 1983, Cawthorne and Cheong 1984, Čerňanská and others 2006, Chartier and others 1998, Díez-Baños and others 2008, Domke and others 2012a,b, Geurden and others 2014, Good and others 2012, Grimshaw and others 1994, Hong and others 1992, 1996, Maingi and others 1996b, Martínez-Valladares and others 2012a, McMahon and others 2013, Mitchell and others 2010, Taylor and others 2009, Traversa and others 2007, Vadlejch and others 2014
	Goats	France, Denmark, Italy, Netherlands, Norway, Spain, UK	Borgsteede and others 1996, Chartier and others 1998, 2001, Domke and others 2012a, Hong and others 1996, Jackson and others 1992, Maingi and others 1996a, Requejo-Fernández and others 1997, Zanzani and others 2014
ML	Sheep	Czech Republic, Denmark, Italy, Netherlands, Slovakia, Spain, Sweden ^b , UK	Alvarez-Sánchez and others 2006, Bartley and others 2004, 2006, Borgsteede and others 1997, Čerňanská and others 2006, Díez-Baños and others 2008, Höglund and others 2009, Maingi and others 1996b, Martínez-Valladares and others 2012a, b, McMahon and others 2013, Taylor and others 2009, Traversa and others 2007, Vadlejch and others 2014
	Goats	Denmark, Italy, Switzerland, UK	Jackson and others 1992, Maingi and others 1996a, Murri and others 2014, Zanzani and others 2014
LEV	Sheep	Denmark, France, Greece, Ireland, Italy, Netherlands, Spain, UK	Alvarez-Sánchez and others 2006, Bartley and others 2004, 2006, Bjørn and others 1991, Borgsteede and others 1997, Chartier and others 1998, Geurden and others 2014, Good and others 2012, Hong and others 1996, Maingi and others 1996b, Martínez-Valladares and others 2012b, McMahon and others 2013, Mitchell and others 2010, Taylor and others 2009, Traversa and others 2007
	Goats	Denmark, France, UK	Chartier and others 2001, Hong and others 1996, Maingi and others 1996a

MDR	Sheep	Denmark, Greece, Ireland, Italy, Netherlands, Spain, UK	Alvarez-Sánchez and others 2006, Bartley and others 2004, 2006, Borgsteede and others 1997, Geurden and others 2014, Good and others 2012, Maingi and others 1996b, Martínez-Valladares and others 2012b, Mitchell and others 2010, Sargison and others 2005, 2007, 2010, Taylor and others 2009, Traversa and others 2007, Wilson & Sargison 2007
	Goats	Denmark, France, UK	Chartier and others 2001, Jackson and others 1992, Maingi and others 1996a

Trichostrongylus spp.

BZ	Sheep	Denmark, France, Greece, Ireland, Italy, Netherlands, Norway, Slovakia, Spain, UK	Alvarez-Sánchez and others 2006, Bjørn and others 1991, Boersema and others 1987, Borgsteede and others 1997, 2007, Čerňanská and others 2006, Chartier and others 1998, Díez-Baños and others 2008, Domke and others 2012a, Geurden and others 2014, Good and others 2012, Maingi and others 1996b, Martínez-Valladares and others 2013, McMahon and others 2013, Mitchell and others 2010, Palcy and others 2010, Taylor and others 2009, Traversa and others 2007
	Goats	Denmark, France, Italy, Netherlands, Norway	Borgsteede and others 1996, Cabaret and others 1995, Chartier and others 1998, 2001, Cringoli and others 2007, Domke and others 2012a, Maingi and others 1996a, Paraud and others 2009, Zanzani and others 2014
ML	Sheep	Denmark, Greece, Italy, Netherlands, Slovakia, Spain, UK	Alvarez-Sánchez and others 2006, Bartley and others 2006, Borgsteede and others 1997, Čerňanská and others 2006, Geurden and others 2014, Maingi and others 1996b, Martínez-Valladares and others 2013, McMahon and others 2013, Traversa and others 2007
	Goats	Denmark, Italy, Switzerland	Artho and others 2007, Maingi and others 1996a, Murri and others 2014, Zanzani and others 2014
LEV	Sheep	Denmark, France, Greece, Ireland, Italy, Netherlands, Spain, UK	Alvarez-Sánchez and others 2006, Bjørn and others 1991, Borgsteede and others 1997, Chartier and others 1998, Geurden and others 2014, Good and others 2012, Maingi and others 1996b, Martínez-Valladares and others 2013, McMahon and others 2013, Mitchell and others 2010, Taylor and others 2009, Traversa and others 2007
	Goats	Denmark, France	Chartier and others 2001, Maingi and others 1996a, Paraud and others 2009

MDR	Sheep	Denmark, Germany, Greece, Ireland, Italy, Netherlands, Spain, UK	Alvarez-Sánchez and others 2006, Borgsteede and others 1997, Geurden and others 2014, Good and others 2012, Maingi and others 1996b, Martínez-Valladares and others 2013, Mitchell and others 2010, Traversa and others 2007, Voigt and others 2012
	Goats	Denmark, France	Chartier and others 2001, Maingi and others 1996a
<hr/> <i>Haemonchus contortus</i>			
BZ	Sheep	France, Germany, Greece, Netherlands, Norway, Slovakia, Sweden, Switzerland, UK	Boersema and others 1987, Borgsteede and others 1997, 2007, Borgsteede and Duyn 1989, Cawthorne and Cheong 1984, Čerňanská and others 2006, Domke and others 2012a,b, Gallidis and others 2012, Geurden and others 2014, Grimshaw and others 1994, Hertzberg and others 2000, Höglund and others 2009, Hong and others 1992, Jordi 1980, Meyer 2001, Scheuerle and others 2009
	Goats	France, Netherlands, Switzerland	Borgsteede and others 1996, Cabaret and others 1995, Chartier and others 1998, 2001, Hertzberg and others 2000, Meyer 2001, Schnyder and others 2005
ML	Sheep	Czech Republic, Germany, Greece, Italy, Netherlands, Slovakia, Switzerland	Artho and others 2007, Borgsteede and others 1997, 2007, Čerňanská and others 2006, Geurden and others 2014, Scheuerle and others 2009, Vadlejch and others 2014, Zanzani and others 2014
	Goats	Germany, Switzerland	Artho and others 2007, Murri and others 2014, Scheuerle and others 2009, Schnyder and others 2005
LEV	Sheep	Greece, Netherlands	Borgsteede and others 1997, Geurden and others 2014
	Goats	France	Chartier and others 2001
MDR	Sheep	Greece, Netherlands	Borgsteede and others 1997, Geurden and others 2014, Van den Brom and others 2013
	Goats	France, Switzerland	Chartier and others 2001, Schnyder and others 2005

Ostertagia ostertagi

BZ	Cattle	Belgium ^b	Borgsteede and others 1992
-----------	---------------	----------------------	----------------------------

ML	Cattle	Germany, Sweden	Demeler and others 2009
-----------	---------------	--------------------	-------------------------

Cooperia oncophora

ML	Cattle	Belgium, Germany, Ireland, Sweden, UK	Areskog and others 2013, Bartley and others 2012, Coles and others 1998, Demeler and others 2009, El-Abdellati and others 2010a,b, McArthur and others 2011, O'Shaughnessy and others 2014, Stafford & Coles 1999
-----------	---------------	--	---

Other^c

BZ	Sheep	Ireland, Netherlands, Slovakia, Spain, UK	Bartley and others 2003, Borgsteede 1986, Burgess and others 2012, Calvete and others 2012, Grimshaw and others 1994, Keane and others 2014, Praslička and others 1994, Várady and others 2006, de Waal, T. and others unpublished observations
-----------	--------------	--	---

LEV	Sheep	Ireland, UK	Burgess and others 2012, Grimshaw and others 1994, Keane and others 2014
------------	--------------	-------------	--

ML	Sheep	Ireland	Keane and others 2014
-----------	--------------	---------	-----------------------

MDR	Sheep	UK	Burgess and others 2012
------------	--------------	----	-------------------------

^aAH = anthelmintic, BZ = benzimidazoles (including pro-BZs), ML = macrocyclic lactones, LEV = levamisole, MDR = multiple drug resistance

^bSuspected resistance

^cOther minor species, or species/genera not identified